arXiv:gr-qc/0309043v3 6 Oct 2003

Tachyonic potential in Bianchi type-I universe

P.K.Suresh
School of Physics,University of Hyderabad.Hyderabad 500 046.India.[*]

(Dated: February 7, 2008)

Motivated form recent string theoretic results, a tachyonic potential is constructed for a spatially homogeneous and anisotropic background cosmology.

PACS numbers: 98.80.Cq

I. INTRODUCTION

Particles with velocities greater than the speed of light, popularly known as tachyon, has been received much attention in cosmology recently [1, 2]. Though the subject of tachyon is still speculative, its various properties has been studied [3]. Recently, Sen [4] has shown that a rolling tachyonic condensate can be studied with the help of the energy density and pressure of an effective fluid. Following these lines, Finstein [5] has obtained an exact solution for a spatially flat isotropic universe in terms of tachyonic potential. Using the tachyonic condensate, Padmanabhan [6]has studied the possibility of expanding universe driven by tachyonic matter. Cosmological inflation driven by the rolling tachyon in the context of the brane world scenario is also investigated [7, 8]. Sami has constructed tachyonic potential which can implements power-law inflation in the brane world cosmology[9]. The possibility of tachyonic inflation with exponential also studied[11]. These studies indicate, the evolution of the tachyon condensate can have cosmological significance and may be useful to understand various cosmological issues. The aforementioned tachyonic condensate has explored in the context of isotropic Friedman-Robertson-Walker (FRW) cosmological model.

The observed large scale isotropy of the universe have been established for times after the era at which the universe become transparent to radiation, and can successfully account by FRW model. This does not mean that the extrapolation to earlier times the model is equally suitable to describe the very early unverse, especially near the Planck scale or string scale. There exist a wide class of anisotropic cosmological models, which also often studying in cosmology, due to various reasons [11]. There are theoretical arguments that sustain the existence of an anisotropic phase that approaches an isotropic case[12]. Also, anisotropic cosmological models are found a suitable candidate to avoid the assumption of specific initial conditions in FRW models. The early universe could also characterized by irregular expansion mechanism. Therefor, it would be useful to explore cosmological models in which anisotropic, existing at early stage of expansion, are damped out in the course of evolution. Interest in such models have been received much attention since 1978[13]. Among the anisotropic models, the Bianchi type - I cosmological model is the simplest one, which is an anisotropic generalization of the Friedman model with Euclidean spatial geometry, and that contains special isotropic cases and allows arbitrary small anisotropy levels [14]. Since present universe is isotropic, it makes an appropriate candidate for studying the possible effects of anisotropy in the early universe on present-day observations. There is evidence that the dynamics of the early universe may have been profoundly influenced by the presence of spatial anisotropie just below the Planck or string scale [15]. Also the solution of the low-energy string cosmological effective action are by their nature anisotropic[16, 17]. Therefor it would be useful to study the role of tachyonic field in Bianchi type -I cosmological model.

Usually, scalar field potential is used to drive the expansion of the universe. Since there exist a mapping between tachyonic and scalar field potentials directly [6], it is reasonable to expect that tachyonic potential can also be used to drive the expansion of the universe. Thus, to study the expansion of the universe, driven by tachyonic source, it is instructive to determine the tachyonic potential in a specific cosmological model under consideration. The present work is to obtain the tachyonic potential in Bianchi type -I unverse.

For the present work we take $c = 1$.

II. PERFECT FLUID AND SCALAR FIELD IN BIANCHI TYPE-I METRIC

Consider spatially homogeneous, anisotropic, and topologically three - torus or Euclidean background metric

$$dS^2 = -dt^2 + \sum_{i=1}^{3} R_i^2(t) \left(dx^i\right)^2 , \qquad (1)$$

where R_1, R_2, and R_3 are the scale factors in x, y, and z directions respectively. The metric (1) is an anisotropic generalization of the Friedman model with Euclidean spatial geometry. The three scale factors are determined vis Einstein equations.

Next, we consider certain procedure to construct some useful relations and can be used to obtain potential for

---

[*] Electronic address: pkssp@uohyd.ernet.in

scalar field and tachyonic field in Bianchi type-I spacetime. For this, assume that the spacetime (1) filled with a perfect fluid. Investigations of Bianchi type-I spacetime filled with a perfect fluid has studied by many authors [18, 19, 20]. The energy -momentum tensor for the perfect fluid can be written as

$$T_{\mu\nu} = -pg_{\mu\nu} + (\varrho + p)v^{\mu}v_{\nu}, \tag{2}$$

where p and ρ are respectively known as pressure and energy density for the perfect fluid. Thus for metric (1) Friedman equations becomes:

$$\frac{\dot{R}_1\dot{R}_2}{R_1R_2} + \frac{\dot{R}_2\dot{R}_3}{R_2R_3} + \frac{\dot{R}_1\dot{R}_3}{R_1R_3} = 8\pi G\rho, \tag{3}$$

$$\begin{aligned} -2\left(\frac{\ddot{R}_1}{R_1} + \frac{\ddot{R}_2}{R_2} + \frac{\ddot{R}_3}{R_3}\right) - \frac{\ddot{R}_1R_2}{R_1R_2} \\ -\frac{\ddot{R}_2R_3}{R_2R_3} - \frac{\ddot{R}_1R_3}{R_1R_3} = 24\pi pG, \end{aligned} \tag{4}$$

Thus

$$\sum_{i=1}^{3}\frac{\ddot{R}_i}{R_i} = -4\pi G\left(\rho + 3p\right). \tag{5}$$

It is convenient to define a parameter

$$\omega(t) \equiv \frac{p\left(t\right)}{\rho\left(t\right)}. \tag{6}$$

The equation of motion for the present situation can be written as

$$d\left(\rho R_1R_2R_3\right) = -\omega\rho d\left(R_1R_2R_3\right), \tag{7}$$

which leads to

$$\frac{\dot{\rho}}{\rho} = -\sum_{i=1}^{3}H_i\left(t\right)\left(\omega\left(t\right) + 1\right). \tag{8}$$

In the anisotropic background metric (1), one can write $\rho \propto (H_1H_2 + H_2H_3 + H_1H_3)$, so that

$$\frac{\dot{\rho}}{\rho} = \frac{\dot{\mathcal{H}}}{\mathcal{H}}, \tag{9}$$

where

$$\mathcal{H} = H_1H_2 + H_2H_3 + H_1H_3. \tag{10}$$

and

$$\begin{aligned} \dot{\mathcal{H}} \ = \ & \dot{H}_1H_2 + \dot{H}_2H_1 + \dot{H}_2H_3 + \\ & \dot{H}_3H_2 + \dot{H}_1H_3 + \dot{H}_3H_1. \end{aligned} \tag{11}$$

Therefor

$$\omega + 1 = -\frac{1}{\sum_{i=1}^{3}H_i}\frac{\dot{\mathcal{H}}}{\mathcal{H}}. \tag{12}$$

The procedure of computing the above relation will be useful for further study.

Since there exist a mapping between the scalar field potential and tachyonic potential [6], it would be useful to compare both potentials in Bianchi type-I cosmological model. Thus, we first obtain the potential for the scalar field.

Consider a $k = 0$ universe with a scalar field as the source and its Lagrangian is given by

$$L = \frac{1}{2}\partial_{\alpha}\varphi\partial^{\alpha}\varphi - U_{\varphi}(\varphi). \tag{13}$$

Therefor the equation of motion for the scalar field for the metric (1)becomes

$$\ddot{\varphi} + \sum_{i=1}^{3}H_i\dot{\varphi} = -\frac{dU_{\varphi}}{d\varphi}. \tag{14}$$

Assume that the evolution of the universe is already specified so that R_i, (i=1,2,3), $H_i = \frac{\dot{R}_i}{R_i}$.. etc, are known functions of time and we need to determine U_{φ}, Since the motion under consideration is spatially homogeneous,i.e., $\varphi(t,x) = \varphi(t)$, the energy density and pressure of the scalar field are respectively given by

$$\rho_{\varphi} = \frac{1}{2}\dot{\varphi} + U_{\varphi}; \qquad p_{\varphi} = \frac{1}{2}\dot{\varphi} - U_{\varphi}. \tag{15}$$

From the definition of ω and (15), it follows that

$$\frac{\dot{\varphi}^2}{U_{\varphi}} = \frac{(1+\omega_{\varphi})}{(1-\omega_{\varphi})} \equiv Q(t), \tag{16}$$

say. Thus $\dot{\varphi}^2 = 2QU_{\varphi}$, differentiating with respect to time and using (12) we get

$$\frac{\dot{U}_{\varphi}}{U_{\varphi}} = -\frac{\dot{Q} + 2Q\sum_{i=1}^{3}H_i}{1+Q}. \tag{17}$$

Integrating this expression and using the definition of $Q(t)$ and adopt the method of obtaining (12) for the scalar field, we find

$$U_{\varphi}(t) = \frac{\mathcal{H}}{8\pi G}\left(1 + \frac{1}{2}\frac{\dot{\mathcal{H}}}{\sum_{i=1}^{3}H_i(\mathcal{H})}\right). \tag{18}$$

Substituting back in the expression $\dot{\varphi}^2 = 2QU_{\varphi}$, we find

$$\varphi(t) = \int dt\sqrt{-\frac{\mathcal{H}}{4\pi G}\frac{\left(\frac{\dot{\mathcal{H}}}{\sum_{i=1}^{3}H_i\mathcal{H}}\right)}{\left(2 + \frac{\dot{\mathcal{H}}}{\sum_{i=1}^{3}H_i\mathcal{H}}\right)}\left(1 + \frac{\dot{\mathcal{H}}}{2\sum_{i=1}^{3}H_i\mathcal{H}}\right)} \tag{19}$$

where $\mathcal{H}$ is given by (10). Eqs (18) and (19) completely solve the task of obtaining a potential U_{φ}, which will lead to a given set of R_1, R_2, and R_3. These equations determine $\varphi(t)$ and $U_{\varphi}(t)$ in terms of R_1, R_2, and R_3 thereby implicitly determine U_{φ}.

III. TACHYON FIELD IN BIANCHI TYPE-I METRIC

Our next aim is to obtain the tachyonic potential in the Bianchi type - I universe. The action for tachyonic field coupled with Einstein gravity in the low energy limit is given by [21]

$$S=\int d^4x\sqrt{-g}\left(\frac{R}{16\pi G}-U_\top(\top)\sqrt{1+\partial_\mu\top\partial^\nu\top}\right). \tag{20}$$

Where g is the determinant of the metric $g_{\mu\nu}$ and R its scalar curvature. Since (20) has a potential function $U_\top$, one could expect that cosmological evolution can be obtained with tachyonic field as the source by an approximate potential taken in a convenient form.

The tachyon field under our consideration is assumed as spatially homogeneous, i.e,$\top(x,t)=\top(t)$, then its energy density $\rho_\top$ and pressure $p_\top$ respectively are given by [4]

$$\rho_\top=\frac{U_\top}{\sqrt{1-\dot\top^2}}. \tag{21}$$

and

$$p_\top=-U_\top\sqrt{1-\dot\top^2}. \tag{22}$$

Consider a universe with $k=0$ for the spacetime (1) filled with a spatially homogeneous tachyon field. The equation of motion of tachyon field in the spacetime (1) takes following form

$$\frac{\ddot\top}{1-\dot\top^2}+\sum_{i=1}^{3}\frac{\dot R_i}{R_i}\dot\top+\frac{U_{,\top}}{U_\top}=0. \tag{23}$$

For the tachyonic field with (21) and (22), we get

$$\omega_\top+1=\dot\top^2. \tag{24}$$

Again the method of obtaining (12) can be used to compute $\omega_\top$ and applying the result in (24), we get

$$\dot\top^2=-\frac{\dot{\mathcal{H}}}{\mathcal{H}\sum_{i=1}^{3}H_i}. \tag{25}$$

Thus it follows

$$\top=\int dt\sqrt{-\frac{\dot{\mathcal{H}}}{\mathcal{H}\sum_{i=1}^{3}H_i}}. \tag{26}$$

The Friedmann equation for the tachyon field can be written by using (21) as

$$\frac{\dot R_1\dot R_2}{R_1R_2}+\frac{\dot R_2\dot R_3}{R_2R_3}+\frac{\dot R_1\dot R_3}{R_1R_3}=8\pi G\rho_\top. \tag{27}$$

Thus, we get

$$\begin{aligned}U_\top &= \frac{\mathcal{H}}{8\pi G}\left(1+\frac{\dot{\mathcal{H}}}{\mathcal{H}\sum_{i=1}^{3}H_i}\right)^{1/2} \\ &= (-\omega_\top)^{1/2}\rho_\top.\end{aligned} \tag{28}$$

Equations (26) and (28) completely solve the problem of determine $U_\top$ and $\top(t)$, for a given set of R_1, R_2 and R_3.

Consider a universe with power-law expansion $R_1=t^{n_1}, R_2=t^{n_2}$, and $R_3=t^{n_3}$, thus, from (25), the complete solution can be obtained as

$$\top=\left(\frac{2}{n_1+n_2+n_3}\right)^{1/2}t+\top_0, \tag{29}$$

and

$$U_\top=\left(\frac{n_1n_2+n_2n_3+n_1n_3}{8\pi G}\right)\left(1-\frac{2}{n_1+n_2+n_3}\right)^{1/2}\frac{1}{t^2}. \tag{30}$$

Combining (29) and (30), we find the potential is

$$\begin{aligned}U_\top &= \frac{1}{4\pi G}\left(\frac{n_1n_2+n_2n_3+n_1n_3}{n_1+n_2+n_3}\right) \\ &\quad\times\left(1-\frac{2}{n_1+n_2+n_3}\right)^{1/2}\frac{1}{(\top-\top_0)^2}.\end{aligned} \tag{31}$$

Thus

$$U_\top\propto\frac{1}{\top^2}. \tag{32}$$

The rapid expansion is possible for the potential (31) with large values of n_i, as in the case of isotropic FRW model. Also the potential has reasonable behavior of $U_\top\rightarrow 0$ as $\top\rightarrow\infty$ though its form for small and intermediate values of $\top$ is not supported by string theory. Comparing the potentials between the scalar field (18) and tachyonic field (28) in Bianchi type - I background metric, it shows that the forms of the both potentials exhibit similar behavior.

To see the asymptotic form of the evolution of the solutions found above, assume that the late time behavior of $\top$ is given by

$$\begin{aligned}\top &= \left(\frac{2}{n_1+n_2+n_3}\right)t+\alpha e^{-\beta t} \\ &= \left(\frac{2}{n_1+n_2+n_3}\right)t+\mathcal{O}(e^{-\beta t})\end{aligned} \tag{33}$$

where n_1, n_2, n_3, α and β are constants with $2\leq n_1+n_2+n_3$.

The value of $n_1+n_2+n_3$ is restricted to 2 because $\top(t)$ to grow proportional to t asymptotically with exponentially small corrections. Thus the asymptotically (late times) behavior of the potential can be written as

$$\begin{aligned}U_\top &\approx \frac{1}{4\pi G}\left(\frac{n_1n_2+n_2n_3+n_1n_3}{n_1+n_2+n_3}\right) \\ &\quad\times\left(1-\frac{2}{n_1+n_2+n_3}\right)^{1/2}\top^{-2}\left[1+\mathcal{O}(\frac{e^{-\beta t}}{t})\right].\end{aligned} \tag{34}$$

If we restrict the total value of n_1, n_2 and n_3, such that

$$\sum_{i=1}^{3} n_i \leq 2, \tag{35}$$

then the potential can have values depending on the values of n_i's. The potential give non-zero value only whenever two n are non-zeros. When $R_1 = R_2 = R_3 = R$ and $n_1 = n_2 = n_3 = n$, then the result (30) is consistent with the isotropic case [6].

IV. CONCLUSIONS

In this brief work, we considered a tachyonic field as the source of gravity in the Bianchi type-I background metric and obtained the tachyonic potential. It is noted that for a large values of n_i's, the potential behaves power-law expansion as in the case of isotropic FRW cosmological model. For a given set of scale factors, $R(t)_i$, it is always possible to obtain a potential, U_{T} such that it result in a cosmological evolution. Thus the tachyonic potentials can be also used as a possible candidate to drive the expansion of anisotropically expanding universe, described by the metric(1). The asymptotic nature of the potential is also discussed. We also obtained potential of scalar field in the Bianchi type-I cosmological model. The present study can account for the tachyonic potential for the Bianchi type-I cosmological model and is found consistent with isotropic FRW cosmological model when n_1=n_2=n_3=n.

Acknowledgments

I thank Prof.Bindu A Bambah for useful discussion. I wish to thank the Director,IUCAA,Pune for warm hospitality and library facilities extended to me and acknowledge the Associateship of IUCAA.

---

[1] G.W.Gibbons, hep-th/0204008.
[2] M.Fairbairn and M.H.G.Tytgat, hep-th/0204070.
[3] Recami, Classical Tachyons and Possible Applications, *La Rivista del Nuovo Cimento della Societa Hallana di Fisica* (1986).
[4] A.Sen,hep-th/0203211;hep-th/0203265; hep-th/0204084.
[5] A.Feinstein,hep-th/02024140.
[6] T.Padmanabhan, hep-th/0204150.
[7] S.Mukohyama,Phys.Rev.D**66**,024009 (2002)[hep-th/0204084]
[8] M.C.Bento,O.Bertolami and A.A.Sen, Phys.Rev.D **67**,063511(2003)[hep-th/0208124]
[9] M.Sami, Mod.Phys.Lett.A **18**,691(2003)[hep-th/0205146]
[10] M.Sami,P.Chingangbam and T.Qureshi,Phys.Rev.D**66**,043530 (2002)[hep-th/0205179]
[11] C.W.Misner, K.S.Thorne, and J.A.Wheeler, *Gravitation*(W.H.Freeman: New York).
[12] C.W.Misner, Astrophysics.J **151**,431(1968).
[13] B.L.Hu and L.Parker, Phys.Rev.D **17**,933(1978).
[14] K.C.Jacob, Astrophysics.J **153**,661(1968).
[15] M.A.H.MacCallum, *in General Relativity- An Einstein Centenary Survey*, edited by S.W.Hawking and W.Israel (Cambridge University Press, Cambridge, England (1979).
[16] G.Veneziano, Phys.Lett.B **265**,287(1991).
[17] M.Gasperini and G.Veneziano, Astropart.Phys.**1**,317(1993).
[18] J.D.Barrow, Nature(London),**272**,211(1978).
[19] V.A.Belinskii and I.M.Khalatnikov, Sov.Phys.JETP **36**,591(1978).
[20] O.Heckmann and E.Schucking, *in Gravitation: An Introduction to Current Research*, edited by L.Witten(Wiley, New York,1952).
[21] A.Frolov, L.Kofman and A.Starobinsky,hep-th/0204187.